arXiv:0903.1811v1 [math.AP] 10 Mar 2009

A Liouville comparison principle for solutions of quasilinear differential inequalities

Vasilii V. Kurta

March 10, 2009

Abstract

This work is devoted to the study of a Liouville comparison principle for entire weak solutions of quasilinear differential inequalities of the form $A(u) + |u|^{q-1}u \le A(v) + |v|^{q-1}v$ on $\mathbb{R}^n$, where $n \ge 1$, $q > 0$, and the operator $A(w)$ belongs to a class of the so-called α-monotone operators. Typical examples of such operators are the p-Laplacian and its well-known modifications for $1 < p \le 2$. The results improve and supplement those in [1].

1 Introduction and Definitions

This work is devoted to the study of a Liouville comparison principle for entire weak solutions of quasilinear differential inequalities of the form

$$A(u) + |u|^{q-1}u \le A(v) + |v|^{q-1}v \tag{1}$$

on $\mathbb{R}^n$, where $n \ge 1$, $q > 0$ and the operator $A(w)$ belongs to a class of the so-called α-monotone operators. Typical examples of such operators are the p-Laplacian

$$\Delta_p(w) := \operatorname{div}\left(|\nabla w|^{p-2}\nabla w\right) \tag{2}$$

for $1 < p \le 2$ and its well-known modification (see, e.g., [2, p. 155]),

$$\widetilde{\Delta}_p(w) := \sum_{i=1}^{n} \frac{\partial}{\partial x_i}\left(\left|\frac{\partial w}{\partial x_i}\right|^{p-2}\frac{\partial w}{\partial x_i}\right) \tag{3}$$

for $n \geq 2$ and $1 < p \leq 2$. The results obtained in this work improve and supplement those in [1]. To prove these results we further develop the approach that was proposed to solve similar problems in wide classes of partial differential equations and inequalities in [3] and [4].

In what follows, $n \geq 1$ is a natural number, $q > 0$ and $\alpha > 1$ are real numbers. Let $A(w)$ be a differential operator given formally by the relation

$$A(w) = \sum_{i=1}^{n} \frac{d}{dx_i} A_i\left(x, \nabla w\right). \tag{4}$$

Assume that the functions $A_i\left(x, \xi\right)$, $i = 1, \ldots, n$, satisfy the Carathéodory conditions on $\mathbb{R}^n \times \mathbb{R}^n$. Namely, they are continuous in ξ at almost all $x \in \mathbb{R}^n$ and measurable in x at all $\xi \in \mathbb{R}^n$.

Definition 1 *Let $n \geq 1$ and $\alpha > 1$. The operator $A(w)$, given by (4), is said to be α-monotone if $A_i\left(x, 0\right) = 0$, $i = 1, \ldots, n$, at almost all $x \in \mathbb{R}^n$, and there exists a positive constant $\mathcal{K}$ such that*

$$0 \leq \sum_{i=1}^{n} \left(\xi_i^1 - \xi_i^2\right) \left(A_i\left(x, \xi^1\right) - A_i\left(x, \xi^2\right)\right) \tag{5}$$

and

$$\left(\sum_{i=1}^{n} \left(A_i\left(x, \xi^1\right) - A_i\left(x, \xi^2\right)\right)^2 \right)^{\alpha/2} \leq \mathcal{K} \left(\sum_{i=1}^{n} \left(\xi_i^1 - \xi_i^2\right) \left(A_i\left(x, \xi^1\right) - A_i\left(x, \xi^2\right)\right) \right)^{\alpha-1} \tag{6}$$

hold at all $\xi^1, \xi^2 \in \mathbb{R}^n$ and almost all $x \in \mathbb{R}^n$.

Note that the condition (5) is the well-known monotonicity condition in PDE theory, while the condition (6) is the proper α-monotonicity condition for differential operators, considered first in [4]; see also [5]. Ibidem one can find algebraic inequalities from which it follows immediately that the p-Laplacian Δ_p and its modification $\widetilde{\Delta}_p$, as well as the weighted p-Laplacian

$$\overline{\Delta}_p(w) := \operatorname{div}\left(a\left(x\right) |\nabla w|^{p-2} \nabla w\right), \tag{7}$$

(see, e.g., [6, p. 55]), with any measurable non-negative uniformly bounded function $a(x)$ on $\mathbb{R}^n$, satisfy the α-monotonicity condition for $\alpha = p$ and $1 < p \leq 2$. Note also that the α-monotonicity condition (6) in the case $\xi^2 = 0$ is in turn a special case of the very general growth condition for quasilinear differential operators, considered first in [7].

Definition 2 *Let $n \geq 1$, $\alpha > 1$ and $q > 0$, and let the operator $A(w)$ be α-monotone. By an entire weak solution of the inequality (1) we understand a pair of functions (u, v) measurable on $\mathbb{R}^n$ which belong to the space $W^1_{\alpha,\mathrm{loc}}(\mathbb{R}^n) \cap L_{q,\mathrm{loc}}(\mathbb{R}^n)$ and satisfy the integral inequality*

$$\int\limits_{\mathbb{R}^n} \left[\sum_{i=1}^n \varphi_{x_i} A_i(x, \nabla u) - |u|^{q-1} u \varphi \right] dx \geq \int\limits_{\mathbb{R}^n} \left[\sum_{i=1}^n \varphi_{x_i} A_i(x, \nabla v) - |v|^{q-1} v \varphi \right] dx \tag{8}$$

for every non-negative function $\varphi \in C^\infty(\mathbb{R}^n)$ with compact support.

2 Results

Theorem 1 *Let $n = 1$, $2 \geq \alpha > 1$ and $q > 0$. Let the operator $A(w)$ be α-monotone, and let (u, v) be an entire weak solution of the inequality (1) on $\mathbb{R}^n$ such that $u(x) \geq v(x)$. Then $u(x) = v(x)$ on $\mathbb{R}^n$.*

Theorem 2 *Let $n = 2$, $\alpha = 2$ and $q > 0$. Let the operator $A(w)$ be α-monotone, and let (u, v) be an entire weak solution of the inequality (1) on $\mathbb{R}^n$ such that $u(x) \geq v(x)$. Then $u(x) = v(x)$ on $\mathbb{R}^n$.*

Theorem 3 *Let $n \geq 2$, $2 \geq \alpha > 1$, $n > \alpha$, $\frac{n(\alpha-1)}{n-\alpha} \geq q > \alpha - 1$ and $q \geq 1$. Let the operator $A(w)$ be α-monotone, and let (u, v) be an entire weak solution of the inequality (1) on $\mathbb{R}^n$ such that $u(x) \geq v(x)$. Then $u(x) = v(x)$ on $\mathbb{R}^n$.*

Theorem 4 *Let $n \geq 2$, $2 \geq \alpha > 1$, $n > \alpha$, $q > \frac{n(\alpha-1)}{n-\alpha}$ and $q \geq 1$, and let the operator $A(w)$ be α-monotone. Then there exists no entire weak solution (u, v) of the inequality (1) on $\mathbb{R}^n$ such that $u(x) \geq v(x)$ and the relation*

$$\limsup_{R \to +\infty} R^{-n + \frac{\alpha(q-\nu)}{q-\alpha+1}} \int\limits_{|x|<R} (u - v)^{q-\nu} dx = +\infty \tag{9}$$

holds with any given $\nu \in (0, \alpha - 1]$.

Remark 1 *If, in Theorem 4, $\nu = \alpha - 1$, then the relation (9) has the form*

$$\limsup_{R\to+\infty} R^{-n+\alpha} \int\limits_{|x|<R} (u-v)^{q-\alpha+1} dx = +\infty. \tag{10}$$

To illustrate the sharpness of Theorem 4 we give Example 1.

Example 1 *For $n \geq 2$, $2 \geq \alpha > 1$, $n > \alpha$, $q > \frac{n(\alpha-1)}{n-\alpha}$, $q \geq 1$ and a suitable constant $c > 0$, the pair (u, v) of the functions*

$$u(x) = c(1 + |x|^{\alpha/(\alpha-1)})^{(1-\alpha)/(q-\alpha+1)} \tag{11}$$

and

$$v(x) = 0 \tag{12}$$

is an entire weak solution of the inequality (1) on $\mathbb{R}^n$ with $A(w) = \Delta_\alpha(w)$ or $A(w) = \widetilde{\Delta}_\alpha(w)$, respectively, such that $u(x) \geq v(x)$ and, for any given $\nu \in (0, \alpha - 1]$, the relation

$$\limsup_{R\to+\infty} R^{-n+\frac{\alpha(q-\nu)}{q-\alpha+1}} \int\limits_{|x|<R} (u-v)^{q-\nu} dx = C_1, \tag{13}$$

with C_1 a certain positive constant, holds.

The two following statements are simple corollaries of Theorem 4.

Theorem 5 *Let $n \geq 2$, $2 \geq \alpha > 1$, $n > \alpha$, $q > \frac{n(\alpha-1)}{n-\alpha}$ and $q \geq 1$, and let the operator $A(w)$ be α-monotone. Then, for any given constants $c > 0$ and $\delta > 0$, there exists no entire weak solution (u, v) of the inequality (1) on $\mathbb{R}^n$ such that $u(x) \geq v(x) + c(1 + |x|^{\alpha/(\alpha-1)})^{(1-\alpha)/(q-\alpha+1)+\delta}$.*

Theorem 6 *Let $n \geq 2$, $2 \geq \alpha > 1$, $n > \alpha$, $q > \frac{n(\alpha-1)}{n-\alpha}$ and $q \geq 1$, and let the operator $A(w)$ be α-monotone. Then, for any given constant $c > 0$ there exists no entire weak solution (u, v) of the inequality (1) on $\mathbb{R}^n$ such that $u(x) \geq v(x) + c$.*

Theorem 7 *Let $n \geq 2$, $2 \geq \alpha > 1$, $n > \alpha$ and $1 \geq q > 0$, and let the operator $A(w)$ be α-monotone. Then there exists no entire weak solution (u, v) of the inequality (1) on $\mathbb{R}^n$ such that $u(x) \geq v(x)$ and the relation*

$$\limsup_{R\to+\infty} R^{\alpha-n} \int\limits_{\{|x|<R\}\cap\{x:\, u(x)\neq v(x)\}} (|u|^{q-1}u - |v|^{q-1}v)(u-v)^{1-\alpha} dx = +\infty \tag{14}$$

holds.

Remark 2 *If, in Theorem 7, $q = 1$, then the relation (14) has the form*

$$\limsup_{R\to+\infty} R^{\alpha-n} \int\limits_{\{|x|<R\}\cap\{x\colon u(x)\neq v(x)\}} (u-v)^{2-\alpha}dx = +\infty. \tag{15}$$

To illustrate the sharpness of Theorem 7 we give three examples.

Example 2 *For $n \geq 2$, $2 \geq \alpha > 1$, $n > \alpha$, $\alpha - 1 > q > 0$, $0 < \mu < \frac{n-\alpha}{n}$, $\lambda = (\alpha-1)/(\alpha-1-q)$, and a suitable constant $c > 0$, the pair (u, v) of the functions*

$$u(x) = c(1+|x|^{\alpha/(\alpha-1)})^{\lambda} + (1+|x|^{\alpha/(\alpha-1)})^{-\mu} \tag{16}$$

and

$$v(x) = c(1+|x|^{\alpha/(\alpha-1)})^{\lambda} \tag{17}$$

is an entire weak solution of the inequality (1) on $\mathbb{R}^n$ with $A(w) = \Delta_\alpha(w)$ or $A(w) = \widetilde{\Delta}_\alpha(w)$, respectively, such that $u(x) \geq v(x)$ and the relation

$$\limsup_{R\to+\infty} R^{\alpha-n} \int\limits_{\{|x|<R\}\cap\{x\colon u(x)\neq v(x)\}} (|u|^{q-1}u - |v|^{q-1}v)(u-v)^{1-\alpha}dx = C_2, \tag{18}$$

with C_2 a certain positive constant, holds.

Example 3 *For $n \geq 2$, $2 \geq \alpha > 1$, $n > \alpha$, $\alpha - 1 > q > 0$, $0 < \mu < \frac{n-\alpha}{n}$, $\lambda > (\alpha-1)/(\alpha-1-q)$, and a suitable constant $c > 0$, the pair (u, v) of the functions (16) and (17) is an entire weak solution of the inequality (1) on $\mathbb{R}^n$ with $A(w) = \Delta_\alpha(w)$ or $A(w) = \widetilde{\Delta}_\alpha(w)$, respectively, such that $u(x) \geq v(x)$ and the relation*

$$\limsup_{R\to+\infty} R^{\alpha-n} \int\limits_{\{|x|<R\}\cap\{x\colon u(x)\neq v(x)\}} (|u|^{q-1}u - |v|^{q-1}v)(u-v)^{1-\alpha}dx = 0 \tag{19}$$

holds.

Example 4 *To illustrate the sharpness of Theorem 7 in the case when $1 \geq q > 0$ and $q > \frac{n(\alpha-1)}{n-\alpha}$, we note that for $n \geq 2$, $2 \geq \alpha > 1$, $n > \alpha$, $1 \geq q > 0$, $q > \frac{n(\alpha-1)}{n-\alpha}$, and a suitable constant $c > 0$, the pair (u, v) of the functions (11) and (12) is an entire weak solution of the inequality (1) on $\mathbb{R}^n$ with*

$A(w) = \Delta_\alpha(w)$ or $A(w) = \widetilde{\Delta}_\alpha(w)$, respectively, such that $u(x) \geq v(x)$ and the relation

$$\limsup_{R\to+\infty} R^{\alpha-n} \int\limits_{\{|x|<R\}\cap\{x:\, u(x)\neq v(x)\}} (|u|^{q-1}u - |v|^{q-1}v)(u-v)^{1-\alpha}dx = C_3, \quad (20)$$

with C_3 a certain positive constant, holds.

Remark 3 *If, in Example 4, $q = 1$, then the relation (20) has the form*

$$\limsup_{R\to+\infty} R^{\alpha-n} \int\limits_{\{|x|<R\}} (u-v)^{2-\alpha}dx = C_3. \quad (21)$$

The following statement, which is an immediate corollary of Theorem 7 and Remark 2, supplements the results of Theorem 3 for the case $q = \alpha - 1$, which is possible in that theorem only if $\alpha = 2$ and $q = 1$.

Theorem 8 *Let $n \geq 3$, $\alpha = 2$ and $q = 1$, and let the operator $A(w)$ be α-monotone. Then there exists no entire weak solution (u, v) of the inequality (1) on $\mathbb{R}^n$ such that $u(x) > v(x)$.*

References

[1] V. V. Kurta, *On a Liouville-type comparison principle for solutions of quasilinear elliptic inequalities*, C. R. Math. Acad. Sci. Paris, 336 (2003), no. 11, 897–900.

[2] J.-L. Lions, *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod, Gauthier-Villars, Paris, 1969.

[3] V. A. Kondrat′ev and E. M. Landis, *Semilinear second-order equations with nonnegative characteristic form*, Mat. Zametki 44 (1988), 457–468.

[4] V. V. Kurta, *Some problems of qualitative theory for nonlinear second-order equations*, Doctoral Dissert., Steklov Math. Inst., Moscow, 1994.

[5] V. V. Kurta, *Comparison principle for solutions of parabolic inequalities*, C. R. Acad. Sci. Paris, Série I, 322 (1996), 1175–1180.

[6] J. Heinonen, T. Kilpeläinen and O. Martio, *Nonlinear potential theory of degenerate elliptic equations*, The Clarendon Press, Oxford University Press, New York, 1993.

[7] V. M. Miklyukov, *Capacity and a generalized maximum principle for quasilinear equations of elliptic type*, Dokl. Akad. Nauk SSSR 250 (1980), 1318–1320.

Address:
Vasilii V. Kurta
Mathematical Reviews
416 Fourth Street, P.O. Box 8604
Ann Arbor, Michigan 48107-8604, USA

FAX: + (734) 996-2916

e-mail: vvk@ams.org